



United States
Department of
Agriculture

NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Routt Area, Colorado, Parts of Rio Blanco and Routt Counties



March 18, 2024

Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	12
Map Unit Descriptions.....	12
Routt Area, Colorado, Parts of Rio Blanco and Routt Counties.....	14
50C—Lintim loam, 3 to 12 percent slopes.....	14
88F—Inchau-Jerry complex, 25 to 50 percent slopes.....	15
94—Dorpat-Reddles complex, 30 to 65 percent slopes.....	17
103—Foidel-Rock outcrop complex, 20 to 60 percent slopes.....	19
139—Maciver stony loam, 3 to 25 percent slopes, extremely stony.....	20
Soil Information for All Uses	23
Soil Reports.....	23
Building Site Development.....	23
Dwellings and Small Commercial Buildings.....	23
Soil Physical Properties.....	26
Engineering Properties.....	26
References	34

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Custom Soil Resource Report

MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

 Soil Map Unit Polygons

 Soil Map Unit Lines

 Soil Map Unit Points

Special Point Features

 Blowout

 Borrow Pit

 Clay Spot

 Closed Depression

 Gravel Pit

 Gravelly Spot

 Landfill

 Lava Flow

 Marsh or swamp

 Mine or Quarry

 Miscellaneous Water

 Perennial Water

 Rock Outcrop

 Saline Spot

 Sandy Spot

 Severely Eroded Spot

 Sinkhole

 Slide or Slip

 Sodic Spot

 Spoil Area

 Stony Spot

 Very Stony Spot

 Wet Spot

 Other

 Special Line Features

Water Features

 Streams and Canals

Transportation

 Rails

 Interstate Highways

 US Routes

 Major Roads

 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
Web Soil Survey URL:
Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Routt Area, Colorado, Parts of Rio Blanco and Routt Counties
Survey Area Data: Version 13, Aug 23, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Jul 2, 2021—Aug 25, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background

MAP LEGEND

MAP INFORMATION

imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
50C	Lintim loam, 3 to 12 percent slopes	2.6	5.3%
88F	Inchau-Jerry complex, 25 to 50 percent slopes	3.0	6.2%
94	Dorpat-Reddles complex, 30 to 65 percent slopes	13.2	27.2%
103	Foidel-Rock outcrop complex, 20 to 60 percent slopes	9.9	20.5%
139	Maciver stony loam, 3 to 25 percent slopes, extremely stony	19.8	40.8%
Totals for Area of Interest		48.6	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it

was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Routt Area, Colorado, Parts of Rio Blanco and Routt Counties

50C—Lintim loam, 3 to 12 percent slopes

Map Unit Setting

National map unit symbol: k0g9

Elevation: 6,560 to 8,200 feet

Mean annual precipitation: 20 to 24 inches

Mean annual air temperature: 38 to 41 degrees F

Frost-free period: 30 to 70 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Lintim and similar soils: 80 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lintim

Setting

Landform: Hills

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Slope alluvium derived from shale

Typical profile

A1 - 0 to 5 inches: loam

A2 - 5 to 20 inches: loam

Bt1 - 20 to 30 inches: clay

Bt2 - 30 to 40 inches: clay

BC - 40 to 65 inches: clay

Properties and qualities

Slope: 3 to 12 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.07 to 0.21 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: High (about 10.2 inches)

Interpretive groups

Land capability classification (irrigated): 6c

Land capability classification (nonirrigated): 6c

Hydrologic Soil Group: C

Ecological site: R048AY247CO - Deep Clay Loam

Hydric soil rating: No

Minor Components

Evna

Percent of map unit: 10 percent
Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: R048AY237CO - Stony Loam
Hydric soil rating: No

Venable

Percent of map unit: 5 percent
Landform: Drainageways
Down-slope shape: Linear
Across-slope shape: Concave
Ecological site: R048AY241CO - Mountain Meadow
Hydric soil rating: Yes

Impass

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: R048BY296CO - Claypan
Hydric soil rating: No

88F—Inchau-Jerry complex, 25 to 50 percent slopes

Map Unit Setting

National map unit symbol: k0hs
Elevation: 6,720 to 7,870 feet
Mean annual precipitation: 20 to 24 inches
Mean annual air temperature: 38 to 41 degrees F
Frost-free period: 30 to 70 days
Farmland classification: Not prime farmland

Map Unit Composition

Inchau and similar soils: 55 percent
Jerry and similar soils: 35 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Inchau

Setting

Landform: Hills

Custom Soil Resource Report

Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Convex
Parent material: Colluvium derived from sandstone and shale

Typical profile

A1 - 0 to 3 inches: cobbly loam
A2 - 3 to 10 inches: loam
Bt - 10 to 18 inches: sandy clay loam
C - 18 to 24 inches: very gravelly sandy clay loam
Cr - 24 to 29 inches: bedrock

Properties and qualities

Slope: 25 to 50 percent
Depth to restrictive feature: 20 to 37 inches to paralithic bedrock
Drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Low to moderately high
(0.00 to 0.28 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: Low (about 3.8 inches)

Interpretive groups

Land capability classification (irrigated): 7e
Land capability classification (nonirrigated): 7e
Hydrologic Soil Group: C
Ecological site: R048AY237CO - Stony Loam
Hydric soil rating: No

Description of Jerry

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Colluvium derived from sandstone and shale

Typical profile

A - 0 to 5 inches: gravelly clay loam
Bt1 - 5 to 12 inches: gravelly clay loam
Bt2 - 12 to 32 inches: gravelly clay loam
Btk - 32 to 52 inches: gravelly clay loam
Bk - 52 to 60 inches: gravelly clay loam

Properties and qualities

Slope: 25 to 50 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.21 to 0.71 in/hr)

Custom Soil Resource Report

Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 10 percent
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: Moderate (about 8.6 inches)

Interpretive groups

Land capability classification (irrigated): 7e
Land capability classification (nonirrigated): 7e
Hydrologic Soil Group: C
Ecological site: R048AY247CO - Deep Clay Loam
Hydric soil rating: No

Minor Components

Routtskin

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: R048AY247CO - Deep Clay Loam
Hydric soil rating: No

Lintim

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Concave
Ecological site: R048AY247CO - Deep Clay Loam
Hydric soil rating: No

94—Dorpat-Reddles complex, 30 to 65 percent slopes

Map Unit Setting

National map unit symbol: k0hz
Elevation: 7,540 to 8,860 feet
Mean annual precipitation: 24 to 28 inches
Mean annual air temperature: 37 to 40 degrees F
Frost-free period: 30 to 70 days
Farmland classification: Not prime farmland

Map Unit Composition

Dorpat and similar soils: 70 percent
Reddles and similar soils: 20 percent

Custom Soil Resource Report

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Dorpat

Setting

Landform: Mountain slopes

Landform position (three-dimensional): Mountainflank

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Colluvium derived from sandstone and shale

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

A - 1 to 3 inches: fine sandy loam

E - 3 to 16 inches: sandy loam

Bt1 - 16 to 30 inches: sandy clay loam

Bt2 - 30 to 46 inches: sandy clay loam

Bt3 - 46 to 60 inches: sandy loam

Properties and qualities

Slope: 30 to 65 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.21 to 0.71 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 8.6 inches)

Interpretive groups

Land capability classification (irrigated): 7e

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: C

Ecological site: F048AY918CO - Spruce-Fir Woodland

Hydric soil rating: No

Description of Reddles

Setting

Landform: Mountain slopes

Landform position (three-dimensional): Mountainflank

Down-slope shape: Linear

Across-slope shape: Concave

Parent material: Colluvium derived from sandstone and shale

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

A - 1 to 10 inches: very fine sandy loam

E - 10 to 15 inches: sandy loam

E/B - 15 to 24 inches: sandy loam

E/B - 24 to 28 inches: clay

Bt1 - 28 to 35 inches: clay

Bt2 - 35 to 60 inches: clay

Custom Soil Resource Report

Cr - 60 to 70 inches: bedrock

Properties and qualities

Slope: 30 to 65 percent

Depth to restrictive feature: 57 to 67 inches to paralithic bedrock

Drainage class: Well drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.07 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: High (about 9.1 inches)

Interpretive groups

Land capability classification (irrigated): 7e

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: D

Ecological site: F048AY918CO - Spruce-Fir Woodland

Hydric soil rating: No

Minor Components

Fulvance

Percent of map unit: 10 percent

Landform: Mountain slopes

Landform position (three-dimensional): Mountainflank

Down-slope shape: Linear

Across-slope shape: Convex

Ecological site: F048AY918CO - Spruce-Fir Woodland

Hydric soil rating: No

103—Foidel-Rock outcrop complex, 20 to 60 percent slopes

Map Unit Setting

National map unit symbol: k0j5

Elevation: 6,560 to 8,530 feet

Mean annual precipitation: 20 to 24 inches

Mean annual air temperature: 38 to 41 degrees F

Frost-free period: 30 to 70 days

Farmland classification: Not prime farmland

Map Unit Composition

Foidel and similar soils: 70 percent

Rock outcrop: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Foidel

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Convex
Parent material: Colluvium derived from sandstone and shale

Typical profile

A1 - 0 to 5 inches: sandy loam
A2 - 5 to 10 inches: loam
B/E - 10 to 15 inches: loam
B/E - 15 to 25 inches: clay loam
Bt - 25 to 60 inches: clay loam

Properties and qualities

Slope: 20 to 60 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.21 to 0.71 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: High (about 10.0 inches)

Interpretive groups

Land capability classification (irrigated): 7e
Land capability classification (nonirrigated): 7e
Hydrologic Soil Group: C
Ecological site: R048AY238CO - Brushy Loam
Hydric soil rating: No

Description of Rock Outcrop

Interpretive groups

Land capability classification (irrigated): 8
Land capability classification (nonirrigated): 8
Hydric soil rating: No

139—Maciver stony loam, 3 to 25 percent slopes, extremely stony

Map Unit Setting

National map unit symbol: k0k6
Elevation: 7,050 to 8,530 feet
Mean annual precipitation: 20 to 24 inches

Custom Soil Resource Report

Mean annual air temperature: 38 to 41 degrees F

Frost-free period: 30 to 70 days

Farmland classification: Not prime farmland

Map Unit Composition

Maciver, extremely stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Maciver, Extremely Stony

Setting

Landform: Hills

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Glaciofluvial deposits derived from igneous and sedimentary rock

Typical profile

A - 0 to 5 inches: stony loam

BA - 5 to 11 inches: very gravelly clay loam

Bt - 11 to 21 inches: very gravelly sandy clay loam

Bk - 21 to 34 inches: very cobbly loam

2Bk - 34 to 65 inches: very cobbly sandy clay loam

Properties and qualities

Slope: 3 to 25 percent

Surface area covered with cobbles, stones or boulders: 5.0 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.21 to 0.71 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 35 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 4.4 inches)

Interpretive groups

Land capability classification (irrigated): 6s

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: C

Ecological site: R048BY237CO - Stony Loam

Hydric soil rating: No

Minor Components

Foidel

Percent of map unit: 5 percent

Landform: Hills

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Concave

Custom Soil Resource Report

Ecological site: R048AY238CO - Brushy Loam

Hydric soil rating: No

Lintim

Percent of map unit: 5 percent

Landform: Hills

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Concave

Ecological site: R048AY247CO - Deep Clay Loam

Hydric soil rating: No

Venable

Percent of map unit: 5 percent

Landform: Drainageways

Down-slope shape: Linear

Across-slope shape: Concave

Ecological site: R048AY241CO - Mountain Meadow

Hydric soil rating: Yes

Soil Information for All Uses

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Building Site Development

This folder contains a collection of tabular reports that present soil interpretations related to building site development. The reports (tables) include all selected map units and components for each map unit, limiting features and interpretive ratings. Building site development interpretations are designed to be used as tools for evaluating soil suitability and identifying soil limitations for various construction purposes. As part of the interpretation process, the rating applies to each soil in its described condition and does not consider present land use. Example interpretations can include corrosion of concrete and steel, shallow excavations, dwellings with and without basements, small commercial buildings, local roads and streets, and lawns and landscaping.

Dwellings and Small Commercial Buildings

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. This table shows the degree and kind of soil limitations that affect dwellings and small commercial buildings.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally

cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Information in this table is intended for land use planning, for evaluating land use alternatives, and for planning site investigations prior to design and construction. The information, however, has limitations. For example, estimates and other data generally apply only to that part of the soil between the surface and a depth of 5 to 7 feet. Because of the map scale, small areas of different soils may be included within the mapped areas of a specific soil.

The information is not site specific and does not eliminate the need for onsite investigation of the soils or for testing and analysis by personnel experienced in the design and construction of engineering works.

Government ordinances and regulations that restrict certain land uses or impose specific design criteria were not considered in preparing the information in this table. Local ordinances and regulations should be considered in planning, in site selection, and in design.

Report—Dwellings and Small Commercial Buildings

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. The table shows only the top five limitations for any given soil. The soil may have additional limitations]

Dwellings and Small Commercial Buildings—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties							
Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50C—Lintim loam, 3 to 12 percent slopes							
Lintim	80	Very limited		Very limited		Very limited	
		Shrink-swell	1.00	Shrink-swell	1.00	Shrink-swell	1.00
						Slope	0.14
88F—Inchau-Jerry complex, 25 to 50 percent slopes							
Inchau	55	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
				Depth to soft bedrock	0.90		
Jerry	35	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Shrink-swell	0.92	Shrink-swell	0.85	Shrink-swell	0.92
94—Dorpat-Reddles complex, 30 to 65 percent slopes							
Dorpat	70	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Shrink-swell	0.01			Shrink-swell	0.01
Reddles	20	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Shrink-swell	0.58	Shrink-swell	1.00	Shrink-swell	0.58
103—Foidel-Rock outcrop complex, 20 to 60 percent slopes							
Foidel	70	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Shrink-swell	0.29	Shrink-swell	0.41	Shrink-swell	0.29
Rock outcrop	30	Not rated		Not rated		Not rated	

Dwellings and Small Commercial Buildings—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties							
Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
139—Maciver stony loam, 3 to 25 percent slopes, extremely stony							
Maciver, extremely stony	85	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Large stones	0.02	Large stones	0.02	Large stones	0.02

Soil Physical Properties

This folder contains a collection of tabular reports that present soil physical properties. The reports (tables) include all selected map units and components for each map unit. Soil physical properties are measured or inferred from direct observations in the field or laboratory. Examples of soil physical properties include percent clay, organic matter, saturated hydraulic conductivity, available water capacity, and bulk density.

Engineering Properties

This table gives the engineering classifications and the range of engineering properties for the layers of each soil in the survey area.

Hydrologic soil group is a group of soils having similar runoff potential under similar storm and cover conditions. The criteria for determining Hydrologic soil group is found in the National Engineering Handbook, Chapter 7 issued May 2007(<http://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=17757.wba>). Listing HSGs by soil map unit component and not by soil series is a new concept for the engineers. Past engineering references contained lists of HSGs by soil series. Soil series are continually being defined and redefined, and the list of soil series names changes so frequently as to make the task of maintaining a single national list virtually impossible. Therefore, the criteria is now used to calculate the HSG using the component soil properties and no such national series lists will be maintained. All such references are obsolete and their use should be discontinued. Soil properties that influence runoff potential are those that influence the minimum rate of infiltration for a bare soil after prolonged wetting and when not frozen. These properties are depth to a seasonal high water table, saturated hydraulic conductivity after prolonged wetting, and depth to a layer with a very slow water transmission rate. Changes in soil properties caused by land management or climate changes also cause the hydrologic soil group to change. The influence of ground cover is treated independently. There are four hydrologic soil groups, A, B, C, and D, and three dual groups, A/D, B/D, and C/D. In the dual groups, the first letter is for drained areas and the second letter is for undrained areas.

The four hydrologic soil groups are described in the following paragraphs:

Custom Soil Resource Report

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

Depth to the upper and lower boundaries of each layer is indicated.

Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. "Loam," for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, "gravelly."

Classification of the soils is determined according to the Unified soil classification system (ASTM, 2005) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 2004).

The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection.

If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest.

Percentage of rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in

Custom Soil Resource Report

the field to weight percentage. Three values are provided to identify the expected Low (L), Representative Value (R), and High (H).

Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field. Three values are provided to identify the expected Low (L), Representative Value (R), and High (H).

Liquid limit and *plasticity index* (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. Three values are provided to identify the expected Low (L), Representative Value (R), and High (H).

References:

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Custom Soil Resource Report

Absence of an entry indicates that the data were not estimated. The asterisk '*' denotes the representative texture; other possible textures follow the dash. The criteria for determining the hydrologic soil group for individual soil components is found in the National Engineering Handbook, Chapter 7 issued May 2007(<http://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=17757.wba>). Three values are provided to identify the expected Low (L), Representative Value (R), and High (H).

Engineering Properties—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties														
Map unit symbol and soil name	Pct. of map unit	Hydrologic group	Depth	USDA texture	Classification		Pct Fragments		Percentage passing sieve number—				Liquid limit	Plasticity index
					Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
			<i>In</i>				<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>
50C—Lintim loam, 3 to 12 percent slopes														
Lintim	80	C	0-5	Loam	CL, ML	A-7-6, A-6	0- 0- 0	0- 0- 0	92-95-100	92-95-100	77-84-94	55-62-70	30-37-44	9-13-17
			5-20	Clay loam, loam, silt loam	CL	A-7-6, A-6	0- 0- 0	0- 0- 0	93-96-100	92-95-100	78-86-95	58-65-73	34-41-48	13-17-21
			20-30	Clay loam, silty clay, clay	CH, CL	A-7-6	0- 0- 0	0- 3- 7	92-97-100	92-97-100	78-92-100	63-76-83	46-57-63	25-33-37
			30-40	Clay loam, silty clay, clay	CH, CL	A-7-6	0- 0- 0	0- 0- 0	92-95-100	92-95-100	78-90-100	63-75-83	46-57-63	25-33-37
			40-65	Clay loam, clay	CH, CL	A-7-6	0- 0- 0	0- 3- 7	84-92-100	83-91-100	73-85-98	58-68-80	44-50-56	25-29-33

Custom Soil Resource Report

Engineering Properties—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties														
Map unit symbol and soil name	Pct. of map unit	Hydrologic group	Depth	USDA texture	Classification		Pct Fragments		Percentage passing sieve number—				Liquid limit	Plasticity index
					Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
			<i>In</i>				<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>
88F—Inchau-Jerry complex, 25 to 50 percent slopes														
Inchau	55	C	0-3	Cobbly loam	CL, ML, SC	A-7-6, A-6	0- 0- 0	7-21- 28	83-91-100	82-91-100	68-81-93	49-59-69	30-37-44	9-13-17
			3-10	Loam, clay loam	CL	A-7-6, A-6	0- 0- 0	0- 7- 7	85-92-93	84-92-92	73-83-89	54-62-68	34-40-48	13-16-21
			10-18	Sandy clay loam	CL, SC	A-7-6, A-2-6	0- 0- 0	0- 0- 0	79-93-93	78-92-92	62-80-85	34-48-52	32-41-47	13-19-23
			18-24	Very gravelly sandy clay loam, gravelly sandy clay loam	GC	A-7-6, A-2-6	0- 0- 0	0- 0- 0	40-45-66	38-43-65	32-37-64	17-21-39	30-35-46	13-17-25
			24-29	Bedrock	—	—	—	—	—	—	—	—	—	—
Jerry	35	C	0-5	Gravelly clay loam	CH, CL, GC	A-7-6	0- 0- 0	0- 0- 7	66-72-78	65-71-78	55-65-72	42-51-57	41-49-53	18-23-25
			5-12	Gravelly clay loam, gravelly clay	CH, CL, GC	A-7-6	0- 0- 0	0- 0- 6	67-73-79	66-72-78	59-66-78	46-53-63	45-49-59	25-27-32
			12-32	Gravelly clay loam, gravelly clay	CH, CL, GC	A-7-6	0- 0- 0	0- 0- 6	61-72-79	59-71-78	53-66-77	42-53-63	44-49-57	25-28-33
			32-52	Gravelly clay loam, gravelly clay	CH, CL, GC	A-7-6	0- 0- 0	0- 0- 7	59-71-78	58-70-77	52-65-76	41-52-62	43-47-55	23-26-32
			52-60	Gravelly clay, gravelly clay loam	CH, CL, GC	A-7-6	0- 0- 0	0- 0- 6	60-72-78	58-71-77	52-65-77	41-52-66	42-47-60	23-27-36

Custom Soil Resource Report

Engineering Properties—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties														
Map unit symbol and soil name	Pct. of map unit	Hydrologic group	Depth	USDA texture	Classification		Pct Fragments		Percentage passing sieve number—				Liquid limit	Plasticity index
					Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
			<i>In</i>				<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>
94—Dorpat-Reddles complex, 30 to 65 percent slopes														
Dorpat	70	C	0-1	Slightly decomposed plant material	PT	A-8	0- 0- 0	0- 0- 0	—	—	—	—	—	—
			1-3	Fine sandy loam	SC-SM, SC	A-2-4, A-4, A-6	0- 0- 0	0- 7- 7	84-90-100	83-90-100	73-83-98	30-37-46	23-29-34	6-10-13
			3-16	Gravelly sandy loam, sandy loam	SC-SM, SC	A-2-4, A-4, A-6	0- 0- 0	0- 6- 11	79-89-100	78-89-100	56-69-82	27-36-45	21-26-31	6-10-13
			16-30	Sandy clay loam, gravelly sandy clay loam	CL, SC	A-7-6, A-6	0- 0- 0	0- 9- 16	81-90-100	80-89-100	65-77-91	36-45-55	34-41-47	17-21-25
			30-46	Gravelly sandy clay loam, sandy clay loam	CL, SC	A-2-6, A-7-6, A-6	0- 0- 0	0- 6- 11	77-89-100	76-88-100	61-77-95	34-45-59	30-37-46	13-18-25

Custom Soil Resource Report

Engineering Properties—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties														
Map unit symbol and soil name	Pct. of map unit	Hydrologic group	Depth	USDA texture	Classification		Pct Fragments		Percentage passing sieve number—				Liquid limit	Plasticity index
					Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
			<i>In</i>				<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>
			46-60	Gravelly sandy loam, sandy loam	SC-SM, SC	A-2-4, A-4, A-6	0- 0- 0	0- 6- 11	79-89-1 00	78-89-1 00	56-69-82	27-36-45	21-26-31	6-10-13
Reddles	20	D	0-1	Slightly decomposed plant material	PT	A-8	0- 0- 0	0- 0- 0	—	—	—	—	—	—
			1-10	Very fine sandy loam	SC-SM, CL, ML	A-4, A-6	0- 0- 0	0- 0- 0	86-93-1 00	85-92-1 00	80-92-1 00	45-54-61	25-32-37	6-10-12
			10-15	Sandy loam	SC-SM, SC	A-2-4, A-4, A-6	0- 0- 0	0- 3- 5	85-92-1 00	84-92-1 00	61-71-81	30-37-44	20-26-30	6-10-13
			15-24	Sandy loam	SC-SM, SC	A-2-4, A-4, A-6	0- 0- 0	0- 3- 5	85-92-1 00	84-92-1 00	61-71-81	30-37-44	20-26-30	6-10-13
			24-28	Clay, clay loam	CH, CL	A-7-6	0- 0- 0	0- 3- 5	85-92-1 00	84-92-1 00	74-85-98	59-69-80	46-52-58	25-29-33
			28-35	Clay, clay loam	CH, CL	A-7-6	0- 0- 0	0- 7- 14	85-92-1 00	84-92-1 00	74-85-98	59-69-80	46-51-58	25-28-33
			35-60	Clay	CH	A-7-6	0- 0- 0	0- 3- 6	84-84-92	83-83-92	68-77-92	59-67-83	50-61-72	29-36-44
			60-70	Bedrock	—	—	—	—	—	—	—	—	—	—
103—Foidel-Rock outcrop complex, 20 to 60 percent slopes														
Foidel	70	C	0-5	Sandy loam	SC, SM	A-2-4, A-4, A-6	0- 0- 0	0- 6- 6	85-92-1 00	84-92-1 00	64-71-81	32-37-44	28-32-38	8-10-12
			5-10	Loam	CL, ML	A-7-6, A-6	0- 0- 0	0- 6- 6	84-92-1 00	83-91-1 00	71-81-96	52-61-73	30-35-45	9-12-18
			10-15	Loam	CL-ML, CL	A-4, A-6	0- 0- 0	0- 5- 5	86-93-1 00	85-92-1 00	71-82-92	51-60-69	21-26-30	6-10-13
			15-25	Clay loam, loam	CH, CL	A-7-6, A-6	0- 0- 0	0- 6- 6	85-92-1 00	84-92-1 00	71-84-94	54-65-74	37-45-50	17-22-25
			25-60	Clay loam, loam	CL	A-7-6, A-6	0- 0- 0	0- 4- 14	93-95-1 00	93-95-1 00	78-87-94	60-68-74	36-43-47	17-22-25

Custom Soil Resource Report

Engineering Properties—Routt Area, Colorado, Parts of Rio Blanco and Routt Counties														
Map unit symbol and soil name	Pct. of map unit	Hydrologic group	Depth	USDA texture	Classification		Pct Fragments		Percentage passing sieve number—				Liquid limit	Plasticity index
					Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
			<i>In</i>				<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>	<i>L-R-H</i>
139—Maciver stony loam, 3 to 25 percent slopes, extremely stony														
Maciver, extremely stony	85	C	0-5	Stony loam	GC-GM, CL, SC	A-6	0- 7- 14	7- 7- 14	64-73-82	63-72-82	52-64-73	36-46-53	27-33-35	7-11-12
			5-11	Very gravelly sandy clay loam, very gravelly clay loam	GC	A-2-6, A-7-6	7- 7- 13	7- 7- 13	44-52-59	42-50-57	34-46-55	26-36-44	32-43-50	13-20-24
			11-21	Very cobbly clay loam, very gravelly sandy clay loam	GC	A-2-6, A-2-7	0- 6- 6	8-12- 37	36-39-53	33-36-51	26-31-48	15-19-30	30-38-46	13-19-25
			21-34	Very cobbly loam	GC-GM, CL, SC	A-2-4, A-6	7- 7- 12	29-35-37	47-76-77	45-75-77	36-67-70	26-49-52	23-30-35	6-12-16
			34-65	Extremely cobbly sandy loam, very cobbly sandy clay loam	GP-GM, GC, SC	A-1-a, A-2-6, A-6	6- 6- 16	15-33-39	38-68-70	35-66-69	25-45-58	11-31-44	17-30-37	1-12-18

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelpdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf